




European Centre for Theoretical Studies
in Nuclear Physics and Related Areas



Parton Distribution Amplitudes for heavy-light pseudoscalar mesons

Daniele Binosi


ECT* - Fondazione Bruno Kessler, Italy

Non perturbative QCD 2018

Sevilla, Spain


November 6-9, 2018

Quantum Chromo Dynamics




- **QCD started 10^{-6} secs. after the Big-Bang:**
it is the glue that binds us all, and understanding its dynamics has profound implications
 - **Explain how massless gluons and light quarks are confined**
and bind together to form hadrons...
 - **...thereby explain the origin of ~98% of the mass in the visible Universe**

- **QCD is *likely* a perfect theory**
nothing needs to be added or changed
 - **Validated** over an incredible energy range: $0 \lesssim E \leq 8000 \text{ GeV}$
 - **Unlikely to break down** at any energy scale (asymptotic freedom)
 - **No intrinsic parameters**, just need one observable to define the scale
 $\Lambda_{\text{QCD}} \simeq 200 - 300 \text{ MeV}$ QCD's “standard kilogram”
- **QCD is *a theory* not an effective theory**
- **However, it is innately nonperturbative**
a priori no idea what such theory can produce



QCD: degrees of freedom



- **QCD basic degrees of freedom:**
matter (quarks); gauge (gluons)

$$q_f^i \begin{cases} \text{color} & i = 1, 2, 3 \\ \text{flavor} & f = u, d, s, c, b, t \end{cases}$$

$$A_\mu^a \begin{cases} \text{color} & a = 1, \dots, 8 \\ \text{spin} & \epsilon_\mu^{\pm, 0} \end{cases}$$

- **QCD action:**
encodes all the dynamics

$$S_{\text{QCD}} = \int d^4x (\mathcal{L}_I + \mathcal{L}_{\text{GF+FPG}})$$

$$\mathcal{L}_{\text{GF+FPG}} = s(\bar{c}^a \mathcal{F}^a - \xi/2 \bar{c}^a b^a)$$





$$\mathcal{L}_I = \bar{q}_f^i (i\gamma^\mu D_\mu - m)_{ij} q_f^j - \frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu}$$

- **Gauge fixing + FP ghost:**
BRST exact, does not appear in spectrum


$$D_\mu = \partial_\mu - ig A_\mu^a T^a$$

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$$





- **QCD (self-)interactions**
dictate the theory's behavior



QCD: asymptotic freedom





- **Gluon self-interaction runs**
implies QCD is non perturbative and quark/gluons are confined



- **QCD interaction strength:**
grows with separation between gluons and quarks
- **Typical scale of hadron physics**
 $r \sim 2 \text{ fm}$ $\alpha \sim 0.5$
- **Perturbation theory breaks down at this distance**
QCD is entirely nonperturbative across almost the entire proton's volume

QCD: Schwinger-Dyson equations



- **Understanding the origin of mass in QCD**
is quite likely inseparable from understanding confinement
- **One possible way of addressing this are Schwinger-Dyson eqs**
which are quantum eom of Green's functions
- **SDEs: nonperturbative, covariant, IR/UV, light/heavy quarks; but:**
infinite system of coupled integral equations
 - **Needs reliable truncation schemes**
plus requires a gauge to be chosen (Landau)
- **Concentrate on SDE for 2-point functions** $\text{---}\bullet\text{---}^{-1}$
 - **Three equations to be considered**
quarks, gluons and ghosts

$$\mathcal{L}_{\text{GF+FPG}} = s(\bar{c}^a \mathcal{F}^a - \xi/2 \bar{c}^a b^a)$$
 - **Gauge fixing + FP ghost:**
BRST exact, does not appear in the spectrum
- **Capture two emergent phenomena**
 - Dynamical mass generation
 - Confinement (?)

Intermezzo














“This ‘telephone’ has too many shortcomings to be seriously considered as a means of communication. The device is inherently of no value to us.”
Western Union internal memo



QCD in the IR

- **Dynamical χ SB**
in the matter sector
- **Saturating gluon propagator**
dynamical gluon mass
- **IR massless ghost**
saturating dressing function
- **IR suppressed 3g vertex**
due to ghost-gluon dynamics



PDAs within a DS framework



- **Meson leading twist PDA**

projection of BS wave-function onto the light front

Chang, Cloet, Cobos-Martinez, Roberts, Schmidt, PRL 110 (2013)

$$f_{\text{PS}} \varphi(x, \zeta) = Z_2(\zeta, \Lambda) \text{Tr}_{\text{CD}} \int_{\text{d}q}^{\Lambda} \delta(n \cdot q_+ - x n \cdot P) \gamma_5 (\gamma \cdot n) \chi_{\text{PS}}(q, P)$$

f_{PS}

- **Meson's decay constant**
(normalization of PDAs) $\int_0^1 \text{d}x \phi(x, \zeta) = 1$

$Z_2(\zeta, \Lambda)$

- **Quark wave-function renormalization**
evaluated at ζ (2 GeV)

$\int_{\text{d}q}^{\Lambda}$

- **Poincaré invariant regularization**
of 4-dimensional integral

$\delta(n \cdot q_+ - x n \cdot P)$

- **n light-cone vector; P meson momentum**
 $n \cdot P = -m_{\text{PS}} \quad P^2 = -m_{\text{PS}}^2$

$\chi_{\text{PS}}(q, P)$

- **BS wave-function:** $\chi_{\text{PS}}(q, P) = S_{f_1}(q_+) \Gamma_{\text{PS}}(q, P) S_{f_2}(q_-)$

$$q_+ = q + \eta P$$

$$q_- = q - (1 - \eta)P$$

$\Gamma_{\text{PS}}(q, P)$

- **BS amplitude**
solution of the homogeneous BS equation

$$\Gamma_{\text{PS}}^{\alpha\beta}(k, P) = \int_q \mathcal{K}^{\alpha\alpha'; \beta\beta'}(q_{\pm}, k_{\pm}) [S_{f_1}(q_+) \Gamma_{\text{PS}}^{\alpha\beta}(q, P) S_{f_2}(q_-)]_{\alpha'\beta'}$$

$S_{f_1}(q_+)$

- **Quark propagator**
solution of the gap equation

$$S_f^{-1}(k) = Z_2(-i\gamma \cdot k + m_f) + Z_1 \int_{\text{d}q}^{\Lambda} g^2 D^{\mu\nu}(k - q) \frac{\lambda^a}{2} \gamma_{\mu} S(q) \frac{\lambda^a}{2} \Gamma_{\nu}(q, k)$$

PDAs within a DS framework



- **Meson leading twist PDA**

projection of BS wave-function onto the light front

Chang, Cloet, Cobos-Martinez, Roberts, Schmidt, PRL 110 (2013)

$$f_{\text{PS}} \varphi(x, \zeta) = Z_2(\zeta, \Lambda) \text{Tr}_{\text{CD}} \int_{\text{d}q}^{\Lambda} \delta(n \cdot q_+ - x n \cdot P) \gamma_5 (\gamma \cdot n) \chi_{\text{PS}}(q, P)$$

- **Not accessible**
in Euclidean space

- **Mellin moments**

used for reconstructing the PDA

$$\langle x^m \rangle = \frac{1}{f_{\text{PS}}} Z_2(\zeta; \Lambda) \text{Tr}_{\text{CD}} \int_{\text{d}q}^{\Lambda} \frac{(n \cdot q_+)^m}{(n \cdot P)^{m+1}} \gamma_5 (\gamma \cdot n) \chi_{\text{PS}}(q; P)$$

- **Two possible representations**

for determining the full amplitude

- **Gegenbauer**

$$\varphi(x) = \mathcal{N} x^\alpha (1-x)^\beta$$

- **Gaussian**

$$\varphi(x) = \mathcal{N} [1 - (2x - 1)^2] e^{\mathfrak{a}^2 (1 - (2x - 1)^2) + \mathfrak{b}^2 (2x - 1)}$$

- **Main difference**

end point behavior @ $x \rightarrow 1$

- **Affects the inverse moment** $\int_0^1 \text{d}x \phi(x)/(1-x)$
large Q^2 behavior of form factors


- **Heavy-heavy limit:** $\varphi(x) \rightarrow \delta(1 - x/2)$

- **Heavy-light limit:** $\varphi(x) \rightarrow \delta(1 - x)$ (at least in principle, see later)

- **Highly oscillatory integral**
in Euclidean space

- **Requires damping factor**
and extrapolation

$$1/(1 + r^2 k^2)^m; \quad r^2 \rightarrow 0$$



Propagators and BS amplitudes




- **Use RL truncation**
for gap and BS equation

$$\Gamma_\nu = \gamma_\nu$$

$$\mathcal{K}[S\Gamma_{\text{PS}}S] = -P^{\mu\nu}\mathcal{G}\gamma_\mu[S\Gamma_{\text{PS}}S]\gamma_\nu$$

$$\text{---}\bullet\text{---}^{-1} = \text{---}^{-1} + \text{---}\bullet\text{---}\bullet\text{---}$$



- **For the interaction**
we set

[Qin, Chang, Liu, Roberts and Wilson, PRC 84 \(2011\)](#)

$$\mathcal{I}(k^2) = k^2 \frac{\mathcal{G}_{\text{IR}}(k^2) + \mathcal{G}_{\text{UV}}(k^2)}{4\pi}$$

$$\mathcal{G}_{\text{IR}}(k^2) = \frac{8\pi^2}{\omega^5} \varsigma^3 e^{-k^2/\omega^2}$$


$$\mathcal{G}_{\text{UV}}(k^2) = \frac{96\pi^2}{25} \frac{1 - e^{-k^2/1[\text{GeV}^2]}}{k^2 \log[e^2 - 1 + (1 + k^2/\Lambda^2)^2]}$$

- **Light quark sector (GeV)**

$$\varsigma = 0.8; \quad \omega = 0.5$$

- **Heavy quark sector (GeV)**

$$\varsigma = 0.6; \quad \omega = 0.8$$




Propagators and BS amplitudes



- **Use RL truncation**
for gap and BS equation

$$\Gamma_\nu = \gamma_\nu$$

$$\mathcal{K}[S\Gamma_{\text{PS}}S] = -P^{\mu\nu}\mathcal{G}\gamma_\mu[S\Gamma_{\text{PS}}S]\gamma_\nu$$



- **For the interaction**
we set

[Qin, Chang, Liu, Roberts and Wilson, PRC 84 \(2011\)](#)

$$\mathcal{I}(k^2) = k^2 \frac{\mathcal{G}_{\text{IR}}(k^2) + \mathcal{G}_{\text{UV}}(k^2)}{4\pi}$$

$$\mathcal{G}_{\text{IR}}(k^2) = \frac{8\pi^2}{\omega^5} \varsigma^3 e^{-k^2/\omega^2}$$


$$\mathcal{G}_{\text{UV}}(k^2) = \frac{96\pi^2}{25} \frac{1 - e^{-k^2/1[\text{GeV}^2]}}{k^2 \log[e^2 - 1 + (1 + k^2/\Lambda^2)^2]}$$

- **Light quark sector (GeV)**

$$\varsigma = 0.8; \quad \omega = 0.5$$

- **Heavy quark sector (GeV)**


$$\varsigma = 0.6; \quad \omega = 0.8$$




- **Heavy sector interaction**

overlaps with gauge sector kernel determination

[DB, Chang, Papavassiliou, Roberts, PLB 742 \(2015\)](#)



Propagators and BS amplitudes




- **SDE solutions**

dramatic change in the analytic structure


- **Normal particle**

mass pole on the real axis



- **SDE solutions**

interactions move the mass pole off axis




- **Complex conj. singularities**
characterized by a dynamically generated mass scale

- **SDE singularities**

impacts our ability to find BS solutions

- **Quark propagators**

need to be known on a parabolic region



- **Cauchy th. not valid**

if singularities are inside the contour

- **Maximize parabola apices**

probe bound states up to $M = \Delta_h + \Delta_l$ ($\Delta_h \gg \Delta_l$)

- **Use freedom to unbalance legs**

in BS equation

$$q_+ = q + \eta P$$

$$q_- = q - (1 - \eta)P$$

$$\eta \in [\Delta_h/M, 1 - \Delta_l/M]$$

- **Still, if gap is too big**


no BS solution exists anyway

Heavy-light systems in RL



- **Choose current masses**
for heavy quarks (in GeV): $m_c = 1.25$, $m_b = 4.35$
- **Fix interaction parameters**
to heavy sector values: $\varsigma = 0.6$; $\omega = 0.8$
- **Evaluate properties of η mesons**

	Exp.		lQCD	
	m_i	f_i	m_i	f_i
η_c	2.98	0.272	2.98	0.238
η_b	9.38	0.501	9.39	/





- **Construct an array of fictitious $h\bar{q}$ pseudoscalar mesons**
down to the lowest m_q RL can accommodate
 - **Charm sector:** $m_q \sim 0.4$
 - **Bottom sector:** $m_q \sim 1.3$
- **Estimate PDAs parameters**
use them to (SPM) extrapolate to lower m_q masses

Masses and decay constants





• Charm sector



	This work		Exp.		lQCD	
	m_i	f_i	m_i	f_i	m_i	f_i
D	1.88(6)	0.200(6)	1.87	0.146	1.87	0.147
D_s	1.93(4)	0.218(1)	1.97	0.182/0.187	1.97	0.175

• Bottom sector




	This work		Exp.		lQCD	
	m_i	f_i	m_i	f_i	m_i	f_i
B	5.30(15)	0.313(17)	5.28	0.149	5.28	0.137
B_s	5.36(13)	0.328(11)	5.37	/	5.37	0.162
B_c	6.32(1)	0.373(2)	6.28	/	/	/



Charm sector




• Gegenbauer



• Gaussian




m_q	α	β	a	b
m_s	$2.212^{+0.228}_{-0.184}$	$0.418^{+0.102}_{-0.123}$	$0.442^{+0.025}_{-0.030}$	$1.404^{+0.025}_{-0.025}$
m_u	$2.107^{+0.254}_{-0.168}$	$0.220^{+0.107}_{-0.143}$	$0.262^{+0.025}_{-0.030}$	$1.436^{+0.025}_{-0.025}$





Charm sector








m_q	α	β	a	b
m_s	$2.212_{-0.184}^{+0.228}$	$0.418_{-0.123}^{+0.102}$	$0.442_{-0.030}^{+0.025}$	$1.404_{-0.025}^{+0.025}$
m_u	$2.107_{-0.168}^{+0.254}$	$0.220_{-0.143}^{+0.107}$	$0.262_{-0.030}^{+0.025}$	$1.436_{-0.025}^{+0.025}$



• Gegenbauer






• Gaussian



Bottom sector






• Gegenbauer



m_q	α	β	$\mathbf{a}$	$\mathbf{b}$
m_c	$6.179^{0.398}_{0.379}$	$1.994^{0.342}_{0.141}$	$1.901^{0.070}_{0.070}$	$2.163^{0.060}_{0.060}$
m_s	$4.660^{0.760}_{1.240}$	$0.230^{0.125}_{0.108}$	$0.621^{0.070}_{0.070}$	$2.174^{0.060}_{0.060}$
m_u	$4.612^{0.714}_{1.287}$	$0.144^{0.054}_{0.118}$	$0.495^{0.070}_{0.070}$	$2.166^{0.060}_{0.060}$

• Gaussian





Bottom sector








m_q	α	β	a	b
m_c	$6.179^{+0.398}_{-0.379}$	$1.994^{+0.342}_{-0.141}$	$1.901^{+0.070}_{-0.070}$	$2.163^{+0.060}_{-0.060}$
m_s	$4.660^{+0.760}_{-1.240}$	$0.230^{+0.125}_{-0.108}$	$0.621^{+0.070}_{-0.070}$	$2.174^{+0.060}_{-0.060}$
m_u	$4.612^{+0.714}_{-1.287}$	$0.144^{+0.054}_{-0.118}$	$0.495^{+0.070}_{-0.070}$	$2.166^{+0.060}_{-0.060}$



• Gegenbauer






• Gaussian



Infinite heavy mass limit




- **What happens when $m_h \rightarrow \infty$?**
on general ground limiting moments saturate
- **Two body problem**
mass of the heavy-object controls overall translational motion of entire system; no effect on the relative motion of the light object
- $\langle 2x - 1 \rangle$ **moment**
measure relative heavy/light momentum
- **Check with relevant moments**
add extra point at intermediate heavy mass





- **Fitting function**

$$a + \frac{b}{m_h}$$

Limiting PDA



- **Limiting PDA @ $m_h \rightarrow \infty$**
can be reconstructed from moments/parameters
- **PDA stable**
against fitting different moms/pars



- **What does a limiting PDA imply?**

Gegenbauer or Gaussian?




- **Main difference**

end point behavior @ $x \rightarrow 1$

- **Affects the inverse moment** $\int_0^1 dx \phi(x)/(1-x)$
large Q^2 behavior of form factors

- **Large factor expected**
up to three times (bottom sector)



- **How to distinguish?**

Use Bayesian determination





[Gao, Chang, Liu PLB 770 \(2017\)](#)

- **Model independent**
determination of the PDA through

$$\Phi(k, P) = \int_{-1}^1 dz \int_0^\infty d\gamma \frac{g(\gamma, z)}{(k^2 + zk \cdot P + \frac{1}{4}P^2 + \Lambda^2 + \gamma)^3}$$

$$\varphi(x) = \mathcal{N} \int_0^\infty d\gamma \frac{g(\gamma, 1-2x)}{\gamma + \Lambda^2 - x(x-1)M^2}$$

- **Use MEM**
to extract g from numerical data of Φ
- **Gegenbauer preferred**
@ end point $x \rightarrow 1$





European Centre for Theoretical Studies
in Nuclear Physics and Related Areas



thank**you**